arXiv:0811.1325v1 [math.GT] 9 Nov 2008

ASYMPTOTIC DIMENSION OF PROPER CAT(0) SPACES WHICH ARE HOMEOMORPHIC TO THE PLANE

NAOTSUGU CHINEN* AND TETSUYA HOSAKA

Abstract. In this paper, we investigate a proper CAT(0) space (X, d) which is homeomorphic to $\mathbb{R}^2$ and we show that the asymptotic dimension $\operatorname{asdim}(X, d)$ is equal to 2.

1. Introduction and preliminaries

In this paper, we study asymptotic dimension of proper CAT(0) spaces which are homeomorphic to $\mathbb{R}^2$.

A metric space (X, d) is *proper* if all closed, bounded sets in (X, d) are compact. We say that a metric space (X, d) is a *geodesic space* if for any $x, y \in X$, there exists an isometric embedding $\xi : [0, d(x, y)] \to X$ such that $\xi(0) = x$ and $\xi(d(x, y)) = y$ (such a ξ is called a *geodesic*).

Let (X, d) be a geodesic space and let T be a geodesic triangle in X. A *comparison triangle* for T is a geodesic triangle $\overline{T}$ in the Euclidean plane $\mathbb{R}^2$ with same edge lengths as T. Choose two points x and y in T. Let $\bar{x}$ and $\bar{y}$ denote the corresponding points in $\overline{T}$. Then the inequality

$$d(x, y) \leq d_{\mathbb{R}^2}(\bar{x}, \bar{y})$$

is called the *CAT(0)-inequality*, where $d_{\mathbb{R}^2}$ is the usual metric on $\mathbb{R}^2$. A geodesic space X is called a *CAT(0) space* if the CAT(0)-inequality holds for all geodesic triangles T and for all choices of two points x and y in T. Details of CAT(0) spaces are found in [1].

In Section 2, we first investigate proper CAT(0) spaces which are homeomorphic to $\mathbb{R}^2$ and we show the following.

Proposition 1.1. *Let (X, d) be a proper* CAT(0) *space which is homeomorphic to $\mathbb{R}^2$. Then, $S(x, r)$ is homeomorphic to $\mathbb{S}^1$ for all $x \in X$ and all $r > 0$. Hence the boundary ∂X is homeomorphic to a circle $\mathbb{S}^1$.*

Let (X, d) be a metric space and let $\mathcal{U}$ be a family of subsets of (X, d). The family $\mathcal{U}$ is said to be *uniformly bounded* if there exists a positive number K such that $\operatorname{diam} U \leq K$ for all $U \in \mathcal{U}$. The family $\mathcal{U}$ is said to be *r-disjoint* if $d(U, U') > r$ for any $U, U' \in \mathcal{U}$ with $U \neq U'$.

2000 *Mathematics Subject Classification.* Primary 20F69; Secondary 54F45, 20F65.

Key words and phrases. asymptotic dimension; CAT(0) space; plane.

*Partly supported by the Grant-in-Aid for Scientific Research (C), Japan Society for the Promotion of Science, No. 19540108.

The *asymptotic dimension* of a metric space (X, d) does not exceed n and we write $\operatorname{asdim}(X, d) \leq n$ if for every $r > 0$ there exist uniformly bounded, r-disjoint families $\mathcal{U}^0, \mathcal{U}^1, \ldots, \mathcal{U}^n$ of subsets of X such that $\bigcup_{k=0}^{n} \mathcal{U}^k$ covers X. The *asymptotic dimension* of a metric space (X, d) is equal to n and we write $\operatorname{asdim}(X, d) = n$ if $\operatorname{asdim}(X, d) \leq n$ and $\operatorname{asdim}(X, d) \not\leq n - 1$.

Asymptotic dimension of groups relates to the Novikov conjecture and there are some interesting recent research on asymptotic dimension (cf. [2], [6], [7], [11], [17]). In [11], Gromov remarks that word hyperbolic groups have finite asymptotic dimension and Roe gave detail of the proof in [14]. Asymptotic dimension of CAT(0) groups and CAT(0) spaces is unknown in general.

The purpose of this paper is to prove the following theorem.

Theorem 1.2. *If* (X, d) *is a proper* CAT(0) *space which is homeomorphic to* $\mathbb{R}^2$*, then* $\operatorname{asdim}(X, d) = 2$*.*

We note that the proper CAT(0) space (X, d) in this theorem need not have an action of some group. We give an example in Section 4.

2. Proper CAT(0) spaces which are homeomorphic to $\mathbb{R}^2$

We first give notation used in this paper.

Notation 2.1. Let the set of all natural number, real number and $[0, \infty)$ denote by $\mathbb{N}$, $\mathbb{R}$ and $\mathbb{R}_+$, respectively. Set $\mathbb{R}^n_+ = \mathbb{R}^{n-1} \times \mathbb{R}_+$, $\mathbb{B}^n = \{x \in \mathbb{R}^n : \sum_{i=1}^{n} x_i^2 \leq 1\}$ and $\mathbb{S}^n = \{x \in \mathbb{R}^{n+1} : \sum_{i=1}^{n+1} x_i^2 = 1\}$. Let Y be a subspace of a metric space (X, d). The interior and the closure of Y in a space X will be denoted by $\operatorname{Int}_X Y$ and $\operatorname{Cl}_X Y$, respectively. Also set $B(x, r) = \{y \in X : d(x, y) \leq r\}$ and $S(x, r) = \{y \in X : d(x, y) = r\}$. We denote the geodesic from x to y in a CAT(0) space (X, d) by $[x, y]$ (cf. [1, Proposition II 1.4]). Set $[x, y) = [x, y] \setminus \{y\}$, $(x, y] = [x, y] \setminus \{x\}$ and $(x, y) = [x, y] \setminus \{x, y\}$.

The following lemma is known.

Lemma 2.2. *Let* (X, d) *be a proper* CAT(0) *space,* $r > 0$ *and* $x_0 \in X$*. Then, the following are satisfied:*

1. $B(x_0, r)$ *is a convex set.*
2. $x_0 \notin [x, y] \subset B(x_0, r)$ *and* $(x, y) \subset B(x_0, r) \setminus S(x_0, r)$ *for any* $x, y \in S(x_0, r)$ *with* $d(x, y) < 2r$*.*
3. (*cf.*[1, Lemma II 5.8 and Proposition II 5.12]) *If* X *is a manifold, for each* $x \in X \setminus \{x_0\}$ *there exists a geodesic line* $\xi : \mathbb{R} \to X$ *such that* $\xi(0) = x_0$ *and* $\xi(d(x_0, x)) = x$*.*

We investigate a proper CAT(0) space which is homeomorphic to $\mathbb{R}^2$.

Notation 2.3. Let $(X, d), r, x_0, x$, and y be as in Lemma 2.2(2). Suppose that X is homeomorphic to $\mathbb{R}^2$. By Lemma 2.2, there exist two geodesic rays $\xi_{x_0,x}, \xi_{x_0,y} : \mathbb{R}_+ \to X$ such that $\xi_{x_0,x}(0) = \xi_{x_0,y}(0) = x_0$, $\xi_{x_0,x}(r) = x$ and $\xi_{x_0,y}(r) = y$. By Lemma 2.2, $\xi_{x_0,x}([r, \infty)) \cup [x, y] \cup \xi_{x_0,y}([r, \infty))$ is homeomorphic to $\mathbb{R}$. Since X is homeomorphic to $\mathbb{R}^2$, by Schönflies Theorem, there exists the component C of $X \setminus \xi_{x_0,x}([r, \infty)) \cup [x, y] \cup \xi_{x_0,y}([r, \infty))$ such that $x_0 \notin C$. Set $\ell(x, y) = S(x_0, r) \cap \mathrm{Cl}_X C$.

We show some lemmas.

Lemma 2.4. *Let (X, d) be a proper $\mathrm{CAT}(0)$ space which is homeomorphic to $\mathbb{R}^2$. Then, $S(x, r)$ is a continuum for all $x \in X$ and all $r > 0$.*

Proof. Let $x_0 \in X$ and $r > 0$. Since $B(x_0, r)$ is a convex set, by duality (cf. [15]),

$$\bar{H}_0(X \setminus B(x_0, r)) \cong \check{H}^1(B(x_0, r)) = 0,$$

thus, $X \setminus \mathrm{Int}_X B(x_0, r) = \mathrm{Cl}_X(X \setminus B(x_0, r))$ is connected. Since there exists a deformation retraction of $X \setminus \mathrm{Int}_X B(x_0, r)$ onto $S(x_0, r)$, $S(x_0, r)$ is connected. □

Lemma 2.5. *Let (X, d) be a proper $\mathrm{CAT}(0)$ space which is homeomorphic to $\mathbb{R}^2$, $r > 0$, $x_0 \in X$, $x, y \in S(x_0, r)$ with $0 < d(x, y) < 2r$ and $z \in \ell(x, y)$. Then*

1. *$\ell(x, y)$ is a continuum,*
2. *$[x, y] \cap [x_0, z] \neq \emptyset$ and*
3. *$d(x, z) \leq d(x, y)$.*

Proof. (1) By Notation 2.3, there exists the component D of $X \setminus R$ such that $C \subset D$, where $R = \xi_{x_0,x}(\mathbb{R}_+) \cup \xi_{x_0,y}(\mathbb{R}_+)$. Since X is homeomorphic to $\mathbb{R}^2$, by Schönflies Theorem, $\mathrm{Cl}_X D$ is homeomorphic to $\mathbb{R}^2_+$. Let D' be a copy of D. Define an equivalent relation $\sim$ in $D \cup D'$ as follows: for $a \in D$ and $a' \in D'$, $a \sim a'$ if and only if $a = a'$, $a \in R$ and $a' \in R'$. Set $B = B(x_0, r) \cap \mathrm{Cl}_X D$, $\widetilde{D} = (D \cup D')/\sim$ and $\widetilde{B} = (B \cup B')/\sim$. Then, there exists a deformation retraction $\mathrm{Cl}_X(D \setminus B)$ onto $\ell(x, y)$, $\widetilde{D}$ is homeomorphic to $\mathbb{R}^2$ and $\widetilde{B}$ is a contractible compact set. By the same method as in the proof of Lemma 2.4, we can show that $\ell(x, y) \cup (\ell(x, y))'/\sim$ is connected. Since there exists the natural surjective map from $\ell(x, y) \cup (\ell(x, y))'/\sim$ onto $\ell(x, y)$, $\ell(x, y)$ is connected.

(2) We may assume that $z \notin \{x, y\}$. By Notation 2.3, there exists the component C of $X \setminus \xi_{x_0,x}([r, \infty)) \cup [x, y] \cup \xi_{x_0,y}([r, \infty))$ such that $x_0 \notin C$ and $z \in C$. Thus, $\xi_{x_0,x}([r, \infty)) \cup [x, y] \cup \xi_{x_0,y}([r, \infty))$ separates between x_0 and z in X. Since $[x_0, z] \subset B(x_0, r)$ is an arc connecting between x_0 and z in X, $[x, y] \cap [x_0, z] \neq \emptyset$.

(3) On the contrary, suppose $d(x, z) > d(x, y)$. By (2), there exists $z' \in [x, y] \cap [x_0, z]$. Since $z' \in [x, y]$,

$$d(x, z') + d(z', y) = d(x, y) < d(x, z) \leq d(x, z') + d(z', z),$$

thus, $d(z', y) < d'(z', z)$. Then,

$$r = d(x_0, y) \leq d(x_0, z') + d(z', y) < d(x_0, z') + d(z', z) = d(x_0, z) = r,$$

a contradiction. $\square$

Lemma 2.6. *Let* (X, d) *be a proper* $\mathrm{CAT}(0)$ *space which is homeomorphic to* $\mathbb{R}^2$, $r, t > 0$, $x_0 \in X$ *and* $y_0 \in S(x_0, r)$. *Then* $S(x_0, r) \cap B(y_0, t)$ *is connected.*

Proof. Set $N = S(x_0, r) \cap B(y_0, t)$. If $t \geq 2r$, $S(x_0, r) \subset B(y_0, t)$. By Lemma 2.4, N is connected. We may assume that $t < 2r$. Take $x \in N$. Since $d(y_0, x) \leq t < 2r$, by Lemma 2.5, we have

$$\ell(y_0, x) \subset S(x_0, r) \cap B(y_0, d(y_0, x)) \subset S(x_0, r) \cap B(y_0, t) = N.$$

Therefore, by Lemma 2.5, $N = \bigcup\{\ell(y_0, x) : x \in N\}$ is connected, which proves the lemma. $\square$

We obtain the following proposition from lemmas above.

Proposition 2.7. *Let* (X, d) *be a proper* $\mathrm{CAT}(0)$ *space which is homeomorphic to* $\mathbb{R}^2$. *Then,* $S(x, r)$ *is homeomorphic to* $\mathbb{S}^1$ *for all* $x \in X$ *and all* $r > 0$.

Proof. By Lemma 2.4 and [16, Theorem 11.21], it suffices to show the following:

(1) $S(x_0, r) \setminus \{y_0, y_1\}$ is non-connected for any $y_0, y_1 \in S(x_0, r)$ with $y_0 \neq y_1$.

(2) $S(x_0, r) \setminus \{y_0\}$ is connected for each $y_0 \in S(x_0, r)$.

We take two points $y_0, y_1 \in S(x_0, r)$ with $y_0 \neq y_1$. By Lemma 2.2, there exist geodesic rays $\xi_{x_0,y_0}, \xi_{x_0,y_1} : \mathbb{R}_+ \to X$ such that $\xi_{x_0,y_0}(0) = \xi_{x_0,y_1}(0) = x_0$, $\xi_{x_0,y_0}(r) = y_0$ and $\xi_{x_0,y_1}(r) = y_1$. By Schönflies Theorem, there exist closed sets Z_0, Z_1 of X such that Z_i is homeomorphic to $\mathbb{R}^2_+$ for $i = 0, 1$, $X = Z_0 \cup Z_1$ and $Z_0 \cap Z_1 \subset \xi_{x_0,y_0}(\mathbb{R}_+) \cup \xi_{x_0,y_1}(\mathbb{R}_+)$ is homeomorphic to $\mathbb{R}$. Since $S(x_0, r) \cap \mathrm{Int}_X Z_i \neq \emptyset$ for $i = 0, 1$, $S(x_0, r) \setminus \{y_0, y_1\}$ is non-connected, which proves (1).

Let $x, y \in S(x_0, r) \setminus \{y_0\}$ with $x \neq y$. By Lemma 2.2, there exist geodesic rays $\xi_{x_0,x}, \xi_{x_0,y} : \mathbb{R}_+ \to X$ such that $\xi_{x_0,x}(0) = \xi_{x_0,y}(0) = x_0$, $\xi_{x_0,x}(r) = x$ and $\xi_{x_0,y}(r) = y$. Set $R = \xi_{x_0,x}(\mathbb{R}_+) \cup \xi_{x_0,y}(\mathbb{R}_+)$. By [1, Proposition 1.4(1), p.160], there exists $z \in [x_0, x)$ such that $\xi_{x_0,x}(\mathbb{R}_+) \cap \xi_{x_0,y}(\mathbb{R}_+) = [x_0, z]$. By Schönflies Theorem, there exists the component C of $X \setminus R$ such that $y_0 \notin C$ and $E_{x,y} = \mathrm{Cl}_X C$ is homeomorphic to $\mathbb{R}^2_+$. Set $L_{x,y} = E_{x,y} \cap S(x_0, r)$. We see $B(x_0, d(x_0, z)) \subset E_{x,y}$ or $B(x_0, d(x_0, z)) \cap E_{x,y} = \{z\}$. Suppose that $B(x_0, d(x_0, z)) \subset E_{x,y}$. We note that $L_{x,y}$, $B(x_0, d(x_0, z))$ and $\{z\}$ are deformation retracts of $\mathrm{Cl}_X(E_{x,y} \setminus B(x_0, r))$, $E_{x,y} \cap B(x_0, r)$ and $B(x_0, d(x_0, z))$, respectively. Thus, $\{z\}$ is a deformation retract of $E_{x,y} \cap B(x_0, r)$. By the same method as in the proof of Lemma 2.5 (1), we can show that $L_{x,y} \cup (L_{x,y})'/\sim$ is a deformation retract of $\mathrm{Cl}_X(E_{x,y} \setminus B(x_0, r)) \cup (\mathrm{Cl}_X(E_{x,y} \setminus B(x_0, r)))'/\sim$ and $\{z\}$ is a deformation retract of $(E_{x,y} \cap B(x_0, r)) \cup (E_{x,y} \cap B(x_0, r))'/\sim$, thus, $L_{x,y}$ is connected. Suppose that $B(x_0, d(x_0, z)) \cap E_{x,y} = \{z\}$. Since $\{z\}$ is a deformation retract of $E_{x,y} \cap B(x_0, r)$, by the same method above, we can show that $L_{x,y}$ is connected.

Fix $y_0' \in S(x_0, r) \setminus \{y_0\}$. Since $S(x_0, r) \setminus \{y_0\} = \bigcup\{L_{x,y_0'} : x \in S(x_0, r) \setminus \{y_0, y_0'\}\}$, it is connected, which proves (2). $\square$

Corollary 2.8. *If (X,d) is a proper* $\mathrm{CAT}(0)$ *space which is homeomorphic to* $\mathbb{R}^2$*, then the boundary* ∂X *of* X *is homeomorphic to* $\mathbb{S}^1$.

We show the following lemma which is used in the proof of the main theorem.

Lemma 2.9. *Let* (X,d) *be a proper* $\mathrm{CAT}(0)$ *space which is homeomorphic to* $\mathbb{R}^2$, $x_0 \in X$, $r,t > 0$ *with* $2t < r$ *and* $x, x' \in S(x_0,r)$ *with* $3t \leq d(x,x') < 2r$. *Then there exist* $y_0, \ldots, y_{3n-1} \in S(x_0,r)$ *and* $m \in \mathbb{N}$ *with* $0 < 3m < 3n-1$ *such that* $y_0 = x$, $y_{3m} = x'$, $t \leq \operatorname{diam} \ell(y_i, y_{i+1}) \leq 2t$, $\{y_0, \ldots, y_{3n-1}\} \cap \ell(y_i, y_{i+1}) = \{y_i, y_{i+1}\}$ *for each* $i = 0, \ldots, 3n-1$, $S(x_0,r) = \ell(y_0,y_1) \cup \cdots \cup \ell(y_{3n-1}, y_{3n})$ *and* $\ell(x,x') = \ell(y_0,y_1) \cup \cdots \cup \ell(y_{3m-1}, y_{3m})$, *where* $y_{3n} = y_0$.

Proof. Set $z_0 = y_0 = x$. By Proposition 2.7, $S(x_0,r)$ is homeomorphic to $\mathbb{S}^1$. Since $S(x_0,r) \not\subset B(z_0,t)$, by Lemma 2.6, $S(x_0,r) \cap B(z_0,t)$, $\ell(x,x')$ and $\ell(x,x') \cap B(z_0,t)$ are arcs. Let z_1 be the end point of $\ell(x,x') \cap B(z_0,t)$ with $z_0 \neq z_1$. By Lemma 2.6, we have $\ell(z_0,z_1) = \ell(x,x') \cap B(z_0,t)$. By Lemma 2.6, $S(x_0,r) \cap B(z_1,t)$ is an arc. Since z_0 and z_1 are the end points of $\ell(z_0,z_1)$ with $d(z_0,z_1) = t$, there exists the end point z_2 of $S(x_0,r) \cap B(z_1,t)$ such that $\operatorname{diam} \ell(z_1,z_2) = t$, $\ell(z_0,z_1) \cap \ell(z_1,z_2) = \{z_1\}$ and $\ell(z_0,z_1) \cup \ell(z_1,z_2) = \ell(x,x') \cap B(z_1,t)$. Thus, by induction, we can take $z_2, \ldots, z_{p+1} \in S(x_0,r)$ and an arc $\ell(z_{i-1},z_i)$ in $\ell(x,x')$ with the end points $\{z_{i-1},z_i\}$ such that $z_{p+1} = x'$, $\ell(z_{i-1},z_i) \cap \ell(z_i,z_{i+1}) = \{z_i\}$ for each $i = 1, \ldots, p$, $\ell(z_{i-1},z_i) \cup \ell(z_i,z_{i+1}) = \ell(x,x') \cap B(z_i,t)$, for each $i = 1, \ldots, p$, $\ell(x,x') = \bigcup_{i=1}^{p+1} \ell(z_{i-1},z_i)$, $\operatorname{diam} \ell(z_{i-1},z_i) = t$ for any $i = 1, \ldots, p$ and $\operatorname{diam} \ell(z_p, z_{p+1}) \leq t$. Let $k \in \mathbb{N}$ and $\delta = 0,1,2$ such that $p = 3k + \delta$. Set $m = k$ and $y_{3m} = z_{p+1}$. If $\delta = 0$, set $y_i = z_i$ for each $i = 1, \ldots, 3m-1$. If $\delta = 1$, set $y_i = z_i$ for each $i = 1, \ldots, 3m-2$ and $y_{3m-1} = z_{p-1}$. If $\delta = 2$, set $y_i = z_i$ for each $i = 1, \ldots, 3(m-1)$, $y_{3m-2} = z_{p-3}$ and $y_{3m-1} = z_{p-1}$. Similarly, we have $y_{3m+1}, \ldots, y_{3n-1} \in \mathrm{Cl}_X(S(x_0,r) \setminus \ell(x,x'))$, which proves the lemma. ☐

3. ASYMPTOTIC DIMENSION OF PROPER $\mathrm{CAT}(0)$ SPACES WHICH ARE HOMEOMORPHIC TO $\mathbb{R}^2$

First we show the following.

Lemma 3.1. *Let* (X,d) *be a proper* $\mathrm{CAT}(0)$ *space which is homeomorphic to* $\mathbb{R}^2$. *Then,* $\operatorname{asdim}(X,d) \geq 2$.

Proof. On the contrary, suppose that $\operatorname{asdim}(X,d) \leq 1$. Let $r > 0$. There exist uniformly bounded, $3r$-disjoint families $\mathcal{U}^0, \mathcal{U}^1$ of subsets of X such that $\mathcal{U}^0 \cup \mathcal{U}^1$ covers X. Since X is homeomorphic to $\mathbb{R}^2$, there exist uniformly bounded, r-disjoint families $\mathcal{V}^0, \mathcal{V}^1$ of subsets of X satisfying the following:

(1) $\mathcal{V}^0 \cup \mathcal{V}^1$ covers X.
(2) Every $V \in \mathcal{V}^0 \cup \mathcal{V}^1$ is a compact topological 2-manifold with boundary.

Let $\varepsilon > 0$ with $\varepsilon < r/2$, let $V \in \mathcal{V}^i$ and let M and M' be two components of V with $d(M, M') = d(M, V \setminus M) < \varepsilon$. Then there exists a disk A in X such that $M \cup A \cup M'$ is connected, $V \cup A$ is a compact topological 2-manifold with boundary and $d(V \cup A, V') > r - \varepsilon$ whenever $V' \in \mathcal{V}^i$ with $V \neq V'$. Thus, we may assume that

(3) $d(M, M') \geq \varepsilon$ for each $V \in \mathcal{V}^0 \cup \mathcal{V}^1$ and each two components M, M' of V.

Since $\mathcal{V}^0 \cup \mathcal{V}^1$ is uniformly bounded, there exists $r \leq s = \sup\{\operatorname{diam} C : C$ is a component of $V \in \mathcal{V}^0 \cup \mathcal{V}^1\} < \infty$. Thus, we may assume that there exists a component C_0 of $V_0 \in \mathcal{V}^0$ such that $s - \varepsilon < \operatorname{diam} C_0 \leq s$. We have $c_0, c_1 \in C_0$ such that $d(c_0, c_1) = \operatorname{diam} C_0$. By Lemma 2.2, there exists a geodesic line $\xi : \mathbb{R} \to X$ such that $\xi(0) = c_0$ and $\xi(\operatorname{diam} C_0) = c_1$. Since $C_0 \cap \xi(\mathbb{R}) \subset \xi([0, \operatorname{diam} C_0])$, there exists the component N of ∂C_0 containing c_0, c_1 which is contained in the closure of the unbounded component of $X \setminus C_0$.

We note that $N \subset \bigcup\{V_1 : V_1 \in \mathcal{V}^1\}$ is homeomorphic to $\mathbb{S}^1$. Since $\mathcal{V}^1$ is r-disjoint, there exists a component C_1 of $V_1 \in \mathcal{V}^1$ such that $N \subset C_1$. Then, there exist $t_0, t_1 \in \mathbb{R}$ with $t_0 < 0 < \operatorname{diam} C_0 < t_1$ such that $\xi(t_0), \xi(t_1) \in C_1 \cap \xi(\mathbb{R}) \subset \xi([t_0, t_1])$. By the similar argument above, we can show there exist a component N' of ∂C_1 containing $\xi(t_0), \xi(t_1)$ and $V_2 \in \mathcal{V}^0$ containing N'. If $V_0 = V_2$, by (3), $d(c_0, \xi(t_0)) \geq \varepsilon$ and $d(c_1, \xi(t_1)) \geq \varepsilon$, i.e., $d(\xi(t_0), \xi(t_1)) > \operatorname{diam} C_0 + 2\varepsilon > s$, which contradicts the definition of s. If $V_0 \neq V_2$, $d(V_0, V_2) > r$. Thus, $d(c_0, \xi(t_0)) > r$ and $d(c_1, \xi(t_1)) > r$, i.e., $d(\xi(t_0), \xi(t_1)) > \operatorname{diam} C_0 + 2r > s$, which contradicts the definition of s. □

We prove the main theorem.

Theorem 3.2. *Let* (X, d) *be a proper* $\operatorname{CAT}(0)$ *space which is homeomorphic to* $\mathbb{R}^2$. *Then,* $\operatorname{asdim}(X, d) = 2$.

Proof. By Lemma 3.1 it suffices to show that $\operatorname{asdim}(X, d) \leq 2$.

Let $r > 0$. Fix $x_0 \in X$ and $k \in \mathbb{N}$ with $k \geq 6$. By Lemma 2.9, there exist $y_{0,0}, \dots, y_{0,3n(0)-1} \in S(x_0, kr)$ such that $2r \leq \operatorname{diam} \ell(y_{0,i}, y_{0,i+1}) \leq 16r$, $\{y_{0,0}, \dots, y_{0,3n(0)-1}\} \cap \ell(y_{0,i}, y_{0,i+1}) = \{y_{0,i}, y_{0,i+1}\}$ for each $i = 0, \dots, 3n(0) - 1$ and $S(x_0, kr) = \ell(y_{0,0}, y_{0,1}) \cup \cdots \cup \ell(y_{0,3n(0)-1}, y_{0,3n(0)})$, where $y_{0,3n(0)} = y_{0,0}$. See Figure 3.2.1. Set $\mathcal{V}_{0,\delta} = \{\ell(y_{0,3i+\delta}, y_{0,3i+1+\delta}) : i = 0, \dots, n(0) - 1\}$ for each $\delta = 0, 1, 2$.



Figure 3.2.1

For every $i = 0, \ldots, 3n(0) - 1$ there exists a geodesic ray $\xi_{0,i} : \mathbb{R}_+ \to X$ such that $\xi_{0,i}(0) = x_0$ and $\xi_{0,i}(kr) = y_{0,i}$. Set $y'_{0,i} = \xi_{0,i}((k+1)r)$ for each $i = 0, \ldots, 3n(0) - 1$. We note $2r \leq d(y_{0,i}, y_{0,i+1}) < d(y'_{0,i}, y'_{0,i+1}) \leq 18r$ for each $i = 0, \ldots, 3n(0) - 1$.

Let $i \in \{0, \ldots, 3n(0) - 1\}$.

Suppose that $d(y'_{0,i}, y'_{0,i+1}) < 12r$. We can take $y_{0,i,0} \in \ell(y_{0,i}, y_{0,i+1})$ and $y'_{0,i,0} \in \ell(y'_{0,i}, y'_{0,i+1})$ such that $r \leq d(y_{0,i}, y_{0,i,0}) = d(y_{0,i+1}, y_{0,i,0}) = d(y_{0,i}, y_{0,i+1})/2 < 6r$ and $\{y_{0,i,0}\} = [x_0, y'_{0,i,0}] \cap S(x_0, kr)$. We note $r < d(y'_{0,i}, y'_{0,i,0}), d(y'_{0,i+1}, y'_{0,i,0}) < 8r$.

Suppose that $12r \leq d(y'_{0,i}, y'_{0,i+1})$. We note that $10r \leq d(y_{0,i}, y_{0,i+1})$. There exist $z_{0,i,0}, z_{0,i,1} \in \ell(y_{0,i}, y_{0,i+1})$ and $z'_{0,i,0}, z'_{0,i,1} \in \ell(y'_{0,i}, y'_{0,i+1})$ such that $d(y_{0,i}, z_{0,i,0}) = d(y_{0,i+1}, z_{0,i,1}) = r$ and $\{z_{0,i,j}\} = [x_0, z'_{0,i,j}] \cap S(x_0, kr)$ for $j = 0, 1$. We note that $d(y'_{0,i}, z'_{0,i,0}), d(y'_{0,i+1}, z'_{0,i,1}) \leq 3r$ and $6r \leq d(z'_{0,i,0}, z'_{0,i,1})$. By Lemma 2.9, there exist $y'_{0,i,1}, \ldots, y'_{0,i,3k_{0,i}-1} \in \ell(y'_{0,i,0}, y'_{0,i,3k_{0,i}})$ such that $2r \leq d(y'_{0,i,j}, y'_{0,i,j+1}) \leq 4r$ and $\ell(y'_{0,i,j}, y'_{0,i,j+1}) \cap \{y'_{0,i,0}, \ldots, y'_{0,i,k_{0,i}}\} = \{y'_{0,i,j}, y'_{0,i,j+1}\}$ for each $j = 0, \ldots, 3k_{0,i} - 1$, where $y'_{0,i,0} = z'_{0,i,0}$ and $y'_{0,i,3k_{0,i}} = z'_{0,i,1}$. See Figure 3.2.2.



Figure 3.2.2

Set $Y_1 = \{y'_{0,i,j} : 0 \le i \le 3n(0) - 1$ and $j = 0, \dots, 3k_{0,i}\}$ and $n(1) \in \mathbb{N}$ with $3n(1) - 1 = |Y_1|$. We can renumber $Y_1 = \{y_{1,i} : i = 0, \dots, 3n(1) - 1\}$ such that

$$\{y' \in S(x_0, (k+1)r) : y \in \bigcup \mathcal{V}_{0,1} \cap \bigcup \mathcal{V}_{0,2}\} \subset \bigcup \mathcal{V}_{1,0},$$
$$\{y' \in S(x_0, (k+1)r) : y \in \bigcup \mathcal{V}_{0,0} \cap \bigcup \mathcal{V}_{0,2}\} \subset \bigcup \mathcal{V}_{1,1},$$
$$\{y' \in S(x_0, (k+1)r) : y \in \bigcup \mathcal{V}_{0,0} \cap \bigcup \mathcal{V}_{0,1}\} \subset \bigcup \mathcal{V}_{1,2},$$

and $\ell(y_{1,i}, y_{1,i+1}) \cap Y_1 = \{y_{1,i}, y_{1,i+1}\}$ for each $i = 0, \dots, 3n(1) - 1$, where we let $y_{1,3n(1)} = y_{1,0}$ and $\mathcal{V}_{1,\delta} = \{\ell(y_{1,3i+\delta}, y_{1,3i+1+\delta}) : i = 0, \dots, n(1) - 1\}$ for each $\delta = 0, 1, 2$. We note that $2r \le \operatorname{diam} V \le 16r$ for all $\delta = 0, 1, 2$ and all $V \in \mathcal{V}_{1,\delta}$.

By induction, for every $m \in \mathbb{N}$ with $m \ge 2$ there exists $Y_m = \{y_{m,i} : i = 0, \dots, 3n(m) - 1\} \subset S(x_0, (k+m)r)$ such that

$$\{y' \in S(x_0, (k+m)r) : y \in \bigcup \mathcal{V}_{m-1,1} \cap \bigcup \mathcal{V}_{m-1,2}\} \subset \bigcup \mathcal{V}_{m,0},$$
$$\{y' \in S(x_0, (k+m)r) : y \in \bigcup \mathcal{V}_{m-1,0} \cap \bigcup \mathcal{V}_{m-1,2}\} \subset \bigcup \mathcal{V}_{m,1},$$
$$\{y' \in S(x_0, (k+m)r) : y \in \bigcup \mathcal{V}_{m-1,0} \cap \bigcup \mathcal{V}_{m-1,1}\} \subset \bigcup \mathcal{V}_{m,2},$$

$\ell(y_{m,i}, y_{m,i+1}) \cap Y_m = \{y_{m,i}, y_{m,i+1}\}$ for each $i = 0, \dots, 3n(m) - 1$ and $2r \le \operatorname{diam} V \le 16r$ for all $\delta = 0, 1, 2$ and all $V \in \mathcal{V}_{m,\delta}$, where we let $\mathcal{V}_{m,\delta} = \{\ell(y_{m,3i+\delta}, y_{m,3i+1+\delta}) : i = 0, \dots, n(m) - 1\}$ for each $\delta = 0, 1, 2$.

Set $\overline{V} = \{x \in B(x_0, (k+m+1)r) \setminus \operatorname{Int}_X B(x_0, (k+m)r) : [x_0, x] \cap V \neq \emptyset\}$ for each $V \in \mathcal{V}_{m,\delta}$, $\overline{\mathcal{V}_{m,\delta}} = \{\overline{V} : V \in \mathcal{V}_{m,\delta}\}$ and $\mathcal{W}_\delta = \left\{W : W \text{ is a component of } \bigcup_{m=0}^{\infty} \overline{\mathcal{V}_{m,\delta}}\right\}$ for each $\delta = 0, 1, 2$. By construction, we have the following:

1. For $V \in \mathcal{V}_{m,\delta}$, $\operatorname{diam} \overline{V} \cap S(x_0, (k+m+1)r) < 12r$ if and only if $\overline{V} \cap \bigcup \overline{\mathcal{V}_{m+1,\delta}} = \emptyset$.
2. Let $\mathcal{V}_{m+1}(V) = \{U \in \mathcal{V}_{m+1,\delta} : \overline{V} \cap U \neq \emptyset\}$ for each $V \in \mathcal{V}_{m,\delta}$. Then, $U \subset \overline{V}$ for $U \in \mathcal{V}_{m+1}(V)$.
3. We have $\mathcal{V}_{m+2}(U) = \emptyset$ for each $V \in \mathcal{V}_{m,\delta}$ and each $U \in \mathcal{V}_{m+1}(V)$ because $\operatorname{diam} U < 12r$ by construction.

For every $\delta = 0, 1, 2$ and every $W \in \mathcal{W}_\delta$ we have

$$\begin{aligned}\operatorname{diam} W &\le \sup\{\operatorname{diam} V : V \in \mathcal{V}_{m,\delta} \text{ for } m \ge 0 \text{ and } \delta = 0, 1, 2\} + 4r \\ &\le 16r + 4r = 20r.\end{aligned}$$

Let $V_i, V_j \in \mathcal{V}_{m,\delta}$ with $V_i \neq V_j$. We show that $d(V_i, V_j) \ge r$. On the contrary, suppose that $d(x, y) < r$ for some $x \in V_i$ and some $y \in V_j$. By Lemma 2.6, let $\ell(x, y)$ denote the arc in $S(x_0, (k+m)r) \cap B(x, d(x, y))$ with the end points $\{x, y\}$. By construction, we have $i = 0, \dots, n(m) - 1$ such that $\ell(y_{m,i}, y_{m,i+1}) \subsetneq \ell(x, y)$. However, $r \le \operatorname{diam} \ell(y_{m,i}, y_{m,i+1}) \le \operatorname{diam} \ell(x, y) = d(x, y) < r$, a contradiction.

Let $\overline{V_i}, \overline{V_j} \in \overline{\mathcal{V}_{m,\delta}}$ with $\overline{V_i} \neq \overline{V_j}$. We show that $d(\overline{V_i}, \overline{V_j}) \ge r$. Let $x' \in \overline{V_i}$ and $y' \in \overline{V_j}$. Set $\{x\} = [x_0, x'] \cap V_i$ and $\{y\} = [x_0, y'] \cap V_j$. By above, $r \le d(V_i, V_j) \le d(x, y)$. Let T be the geodesic triangle consisting of three points x_0, x', y', let

$\overline{T}$ be a comparison triangle for T in $\mathbb{R}^2$ and let $\overline{x_0}, \overline{x}, \overline{y}, \overline{x'}$, and $\overline{y'}$ denote the corresponding points in $\overline{T}$. Since X is a CAT(0) space, we have

$$r \leq d(x,y) \leq d_{\mathbb{R}^2}(\overline{x}, \overline{y}) \leq d_{\mathbb{R}^2}(\overline{x'}, \overline{y'}) = d(x', y'),$$

thus, $d(\overline{V_i}, \overline{V_j}) \geq r$.

Let $\overline{V_i} \in \overline{\mathcal{V}_{m,\delta}}$ and $\overline{V_j} \in \overline{\mathcal{V}_{m+1,\delta}}$ with $\overline{V_i} \cap \overline{V_j} = \emptyset$. Set $W_j = \{[x_0, x] \cap S(x_0, (k+m)) : x \in \overline{V_j}\}$. By the definition of $y'_{m,i,j}$'s, similarly, we can show $d(V_i, W_j) \geq r$. Since X is a CAT(0) space, by the same method we can obtain that $d(\overline{V_i}, \overline{V_j}) \geq r$. By (1), (2) and (3), we have $d(W, W') \geq r$ for any $W, W' \in \mathcal{W}_\delta$ with $W \neq W'$.

Let $\mathcal{U}_0 = \{U : U \text{ is a component of } B(x_0, kr) \cup \bigcup \mathcal{W}_0\}$ and $\mathcal{U}_\delta = \mathcal{W}_\delta$ for $\delta = 1, 2$. By above, $\mathcal{U}_0 \cup \mathcal{U}_1 \cup \mathcal{U}_2$ is a uniformly bounded cover of (X, d) and $d(U, U') \geq r$ for any $U, U' \in \mathcal{U}_\delta$ with $U \neq U'$, which proves the theorem. □

4. Application

As an application of Theorem 3.2, we obtain the following corollary.

Corollary 4.1. *Let (W, S) be a Coxeter system. If the boundary $\partial\Sigma(W, S)$ of $\Sigma(W, S)$ is homeomorphic to $\mathbb{S}^1$, then* asdim$W = 2$.

Proof. Let (W, S) be a Coxeter system whose boundary $\partial\Sigma(W, S)$ is homeomorphic to $\mathbb{S}^1$. Then the Coxeter group W is a virtual Poincaré duality group, and for some $\tilde{S} \subset S$,

$$W = W_{\tilde{S}} \times W_{S \setminus \tilde{S}},$$

where the nerve $N(W_{\tilde{S}}, \tilde{S})$ is homeomorphic to $\mathbb{S}^1$ and $W_{S\setminus\tilde{S}}$ is finite ([4], cf. [12]). Then the Davis complex $\Sigma(W, S)$ splits as

$$\Sigma(W, S) = \Sigma(W_{\tilde{S}}, \tilde{S}) \times \Sigma(W_{S\setminus\tilde{S}}, S \setminus \tilde{S}).$$

Here $\Sigma(W_{\tilde{S}}, \tilde{S})$ is homeomorphic to $\mathbb{R}^2$ and $\Sigma(W_{S\setminus\tilde{S}}, S \setminus \tilde{S})$ is bounded. By Theorem 3.2, we obtain that asdim$\Sigma(W, S) = 2$. Hence asdim$W = 2$. □

In general, it is known that every Coxeter group has finite asymptotic dimension ([6], cf. [8]).

Example 4.2. Let $m \in \mathbb{N}$ and let $D_m \subset \mathbb{R}^2$ be a regular m-polygon with a metric $d_m = d_{\mathbb{R}^2}|_{D_m}$ and edges $e_1, \ldots, e_m$ such that $\operatorname{diam} e_i = 1$ for each $i = 1, \ldots, n$. We consider a noncompact cell 2-complex (Σ, d) with a triangulation $\mathcal{T}$ as follows.

1. For every $\sigma \in \mathcal{T} \setminus \mathcal{T}^{(1)}$ there exist $m(\sigma) \in \mathbb{N}$ and a simplicial isometry f_σ from $(D_{m(\sigma)}, d_{m(\sigma)})$ onto $(|\sigma|, d|_{|\sigma|})$.
2. $|\{\sigma \in \mathcal{T} \setminus \mathcal{T}^{(1)} : \tau < \sigma\}| = 2$ for each $\tau \in \mathcal{T}^{(1)} \setminus \mathcal{T}^{(0)}$.
3. $r(v) = \sum\{\pi - 2\pi/m(\sigma) : v < \sigma \in \mathcal{T} \setminus \mathcal{T}^{(1)}\} \geq 2\pi$ for each $v \in \mathcal{T}^{(0)}$.

(4) for any $x, y \in \Sigma$

$$d(x,y) = \min\{\sum_{j=1}^{k} d_{m(\sigma)}(f_{\sigma_j}^{-1}(x_{j-1}), f_{\sigma_j}^{-1}(x_j)) : \\ x = x_0 \in |\sigma_1|, x_j \in |\sigma_j| \cap |\sigma_{j+1}| (1 \leq j < k), y = x_k \in |\sigma_k|\}.$$

By [3], every (Σ, d) above is a CAT(0) space which is homeomorphic to $\mathbb{R}^2$, hence we obtain that $\operatorname{asdim}(\Sigma, d) = 2$ from Theorem 3.2. Here we note that (Σ, d) need not have an action of some group and (Σ, d) is neither a euclidean nor a hyperbolic plane in general.

REFERENCES

[1] M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*, Springer-Verlag, Berlin, 1999.

[2] G. Bell and A. N. Dranishnikov, *Asymptotic dimension*, Topology Appl. 155 (2008), 1265–1296..

[3] M. W. Davis, *Nonpositive curvature and reflection groups*, in Handbook of geometric topology (Edited by R. J. Daverman and R. B. Sher), pp. 373–422, North-Holland, Amsterdam, 2002.

[4] M. W. Davis, *The cohomology of a Coxeter group with group ring coefficients*, Duke Math. J. 91 (1998), 297–314.

[5] A. N. Dranishnikov, *Asymptotic topology*, (Russian) Uspekhi Mat. Nauk 55 (2000), 71–116; translation in Russian Math. Surveys 55 (2000), 1085–1129

[6] A. N. Dranishnikov and T. Januszkiewicz, *Every Coxeter group acts amenably on a compact space*, Proceedings of the 1999 Topology and Dynamics Conference, Salt Lake City, UT, Topology Proc. 24 (1999), 135–141.

[7] A. N. Dranishnikov, J. Keesling and V. V. Uspenskij, *On the Higson corona of uniformly contractible spaces*, Topology 37 (1998), 791-803.

[8] A. N. Dranishnikov and V. Schroeder, *Embedding of hyperbolic Coxeter groups into products of binary trees and aperiodic tilings*, arXiv:math.GR/0504566v1.

[9] R. Engelking, *General Topology*, Helderman Verlag, Berlin, 1989.

[10] R. Engelking, *Theory of Dimensions Finite and Infinite*, Helderman Verlag, Berlin, 1995.

[11] M. Gromov, *Asymptotic invariants for infinite groups*, Geometric Group Theory (G.A. Niblo and M.A. Roller, eds.), LMS Lecture Notes, vol. 182, Cambridge University Press, Cambridge, 1993, pp.1–295.

[12] T. Hosaka, *On the cohomology of Coxeter groups*, J. Pure Appl. Algebra 162 (2001), 291–301.

[13] G. Moussong, *Hyperbolic Coxeter groups*, Ph.D. thesis, Ohio State University, 1988.

[14] J. Roe, *Hyperbolic groups have finite asymptotic dimension*, Proc. Amer. Math. Soc. 133 (2005), 2489-2490.

[15] E. H. Spanier, *Algebraic topology*, McGraw-Hill Book Co., New York-Toronto, Ont.-London 1966

[16] R. L. Wilder, *Topology of Manifolds*, AMS Colloquium Publications, 32. AMS, New York, N. Y., 1949

[17] G. Yu, *The Novikov conjecture for groups with finite asymptotic dimension*, Annals of Math. 147 (1998), no. 2, 325–355.

Okinawa National College of Technology, Nago-shi Okinawa 905-2192, Japan
E-mail address: `chinen@okinawa-ct.ac.jp`

Department of Mathematics, Faculty of Education, Utsunomiya University, Utsunomiya, 321-8505, Japan
E-mail address: `hosaka@cc.utsunomiya-u.ac.jp`